

A Comparison of Two Rapid Biological Assessment Sampling Methods for Macroinvertebrates

Brad Autrey

U.S. EPA Office of Research and Development (ORD), National Exposure Research Laboratory (NERL), Ecological Exposure Research Division (EERD)

(513) 569-7368

autrey.brad@epa.gov

Authors: Brad Autrey¹, Karen Blocksom¹, Margaret Passmore², Lou Reynolds², Joseph Flotemersch¹

¹U.S. EPA ORD/NERL/EERD, Cincinnati, OH

²U.S. EPA/Region 3, Wheeling, WV

Keywords: macroinvertebrates, method comparison, collaboration, biological assessment, decision-making

In 2003, the Office of Research and Development's (ORD's) National Exposure Research Laboratory (NERL) initiated a collaborative research effort with U.S. Environmental Protection Agency's (U.S. EPA) Region 3 to conduct a study comparing two rapid biological assessment methods for collecting stream macroinvertebrates. One method focuses sampling in a single fast-water habitat (riffles), and the other samples multiple habitats within a stream. The single-habitat method has been widely used in the U.S. for biological assessment of streams for over a decade. However, Region 3 biologists recognized that in the Piedmont and Northern Piedmont regions of the U.S., riffle habitat is less abundant in streams. In streams of these regions, the traditionally used single-habitat method was compared with the more recently proposed multiple-habitat sampling method to determine whether the multiple-habitat method is more effective in these streams. The success of this study depended on a strong collaboration between U.S. EPA Region 3 and the ORD to select sites, conduct field sampling, and analyze the data. While the ORD scientists were responsible for careful analysis of those data, Region 3 biologists were instrumental in helping the NERL put research results into a context useful to state agencies. Region 3 was able to involve biologists at the Virginia Department of Environmental Quality in this process, providing valuable feedback on specific aspects of analysis. As a result of the collaboration between the ORD and Region 3, the results of this methods comparison study will have a more immediate impact on decision-making at the state level.

Although this work was reviewed by the U.S. EPA and approved for publication, it may not necessarily reflect official Agency policy.